


S. Stone  
Sept. 2012



# New Physics from Flavour

Sept., 2012

## ■ Dark Matter



Gravitational  
lensing

- Dark Energy: Cosmological constant
- Hierarchy Problem: Divergent quantum corrections to go from Electroweak scale  $\sim 100$  GeV to Planck scale of Energy  $\sim 10^{19}$  GeV without “fine tuning” quantum corrections
- *All of the above may only be related to Gravity*


# Reasons for NP

- Flavor problem: Why 3 replications of quarks & leptons?
- Baryogenesis: The amount of CP Violation observed thus far in the quark sector is too small:  $(n_B - n_{\bar{B}})/n_\gamma \sim 10^{-20}$  but  $\sim 6 \times 10^{-10}$  is needed. Thus New Physics must exist to generate needed CP Violation
- To explain the values of CKM couplings,  $V_{ij}$ , (both neutrino & quark)
- To explain the masses of fundamental objects. Are they related to the  $V_{ij}$ 's?

# Quark Mixing & CKM Matrix

- In SM charge  $-1/3$  quarks (d, s, b) are mixed
- Described by CKM matrix (also  $\nu$  are mixed)

$$V_{\left(\frac{2}{3}, -\frac{1}{3}\right)} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$



$$= \begin{pmatrix} 1 - \lambda^2 / 2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \lambda^2 / 2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + O(\lambda^4)$$



- $\lambda=0.225$ ,  $A=0.8$ , constraints on  $\rho$  &  $\eta$
- These are fundamental constants in SM



# CKM vs. PMNS

CKM


PMNS



Area  $\sim V^2$

Why these values? Are the two related? Are they related to masses?

# Masses



12 orders of magnitude differences not explained;  $t$  quark as heavy as Tungsten

# Theorists task

- A given theoretical model must explain all the data




*Model must thread through all experimental constraints (12 axe handles). One measurement can, in principle, defeat the theorist, but we seek a consistent pattern.*

# Flavor Physics as a NP discovery tool

- While measurements of CKM parameters & masses are fun, the main purpose of Flavor Physics is to find and/or define the properties of physics beyond the SM
- FP probes large mass scales via virtual quantum loops. An example, of the importance of such loops are changes in the W mass

□  $M_W$  changes due to  $m_t$   $\frac{dM_W}{dm_t} \propto \frac{m_t}{M_W}$



□  $M_W$  changes due to  $m_H$   $\frac{dM_W}{dm_H} \propto -\frac{dm_H}{M_H}$




# Ex. of Strong Constraints on NP

## ■ Inclusive $b \rightarrow s \gamma$ , ( $E_\gamma > 1.6$ GeV)

- Measured  $(3.37 \pm 0.23) \times 10^{-4}$
- Theory  $(3.15 \pm 0.23) \times 10^{-4}$  (NNLL) Misiak arXiv:1010.4896
- Ratio =  $1.07 \pm 0.10$ , Limits most NP models
- Example 2HDM  
 $m(H^+) > 385$  GeV



New BaBar  
 $(3.31 \pm 0.35) \times 10^{-4}$   
See G. Eigen's  
ICHEP talk




Misiak et. al hep-ph/0609232,  
See also A. Buras et. al,  
arXiv:1105.5146




# Limits on New Physics


- It is oft said that we have not seen New Physics, yet what we observe is the sum of Standard Model + New Physics. How to set limits on NP?
- One hypothesis: assume that tree level diagrams are dominated by SM and loop diagrams could contain NP



Tree diagram example




Loop diagram example




# What are limits on NP from quark decays?

- Tree diagrams are unlikely to be affected by physics beyond the Standard Model




# CP Violation in $B^0$ & $K^0$ Only

- Absorptive (Imaginary) part of mixing diagram should be sensitive to New Physics. Lets compare





# They are Consistent




- But consistency is only at the 5% level
- Limits on NP can be derived from difference

# Flavor as a High Mass Probe

## ■ Already excluded ranges from box diagrams

□  $\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{SM}} + \frac{c_i}{\Lambda_i^2} O_i$ , take  $c_i \sim 1$



### Ways out

1. New particles have large masses  $\gg 1$  TeV
2. New particles have degenerate masses
3. Mixing angles in new sector are small, same as in SM (MFV)
4. The above already implies strong constraints on NP


Sept., 2012

See: Isidori, Nir  
& Perez arXiv:1002.0900;  
Neubert EPS 2011 talk






# Neutral Meson Mixing

- Neutral mesons can transform into their anti-particles via 2<sup>nd</sup> order weak interactions
- Short distance transition rate depends on
  - mass of intermediate  $q_i$ , the heavier the larger, favors s & b since t is allowed
  - CKM elements  $V_{ij}$




*New particles possible in the loop*



# Mixing & CPV Definitions

- Mixing & Decay:



$$i \frac{d}{dt} \begin{pmatrix} B_s^0 \\ \bar{B}_s^0 \end{pmatrix} = \begin{pmatrix} M_{11} - \Gamma_{11}/2 & M_{12} - i\Gamma_{12}/2 \\ M_{12}^* - i\Gamma_{12}^*/2 & M_{22} - i\Gamma_{22}/2 \end{pmatrix} \begin{pmatrix} B_s^0 \\ \bar{B}_s^0 \end{pmatrix}$$

- $|M_L\rangle = p|M^0\rangle + q|\bar{M}^0\rangle$ ,  $|M_H\rangle = p|M^0\rangle - q|\bar{M}^0\rangle$ ,
- $m_{B_s} = (M_H + M_L)/2$ ,  $\Delta M = M_H - M_L$ ,  
 $1/\tau_{B_s} = \Gamma = (\Gamma_H + \Gamma_L)/2$ ,  $\Delta\Gamma = \Gamma_L - \Gamma_H$ ,
- $y \equiv \Delta\Gamma/2\Gamma$

# CPV Time Evolution

- Consider CP eigenstate  $a[f(t)] = \frac{\Gamma(\bar{M} \rightarrow f) - \Gamma(M \rightarrow f)}{\Gamma(\bar{M} \rightarrow f) + \Gamma(M \rightarrow f)}$  where  $f$  is a
- Define  $A_f \equiv A(M \rightarrow f)$ ,  $\bar{A}_f \equiv A(\bar{M} \rightarrow f)$ ,  $\lambda_f = \frac{p}{q} \frac{\bar{A}_f}{A_f}$
- $\lambda_f$  is a function of  $V_{ij}$  in SM

$$\Gamma(M \rightarrow f) = N_f |A_f|^2 e^{-\Gamma t} \left( \cosh \frac{\Delta\Gamma t}{2} - \text{Re } \lambda_f \sinh \frac{\Delta\Gamma t}{2} - \text{Im } \lambda_f \sin(\Delta M t) \right)$$



$$\Gamma(\bar{M} \rightarrow f) = N_f |A_f|^2 e^{-\Gamma t} \left( \cosh \frac{\Delta\Gamma t}{2} - \text{Re } \lambda_f \sinh \frac{\Delta\Gamma t}{2} + \text{Im } \lambda_f \sin(\Delta M t) \right)$$

See Nierste  
arXiv:0904.1869 [hep-ph]

# CPV in $B_s \rightarrow J/\psi X$

- Interference between mixing & decay

- For  $f = J/\psi \phi$  or  $J/\psi \pi^+ \pi^-$




$$\varphi_s^{SM} \equiv -2\beta_s = -2 \arg \left( -\frac{V_{ts} V_{tb}^*}{V_{cs} V_{cb}^*} \right) = -0.04 \text{ rad}$$

- Small CPV expected, good place for NP to appear
- $B_s \rightarrow J/\psi \phi$  is not a CP eigenstate, as it's a vector-vector final state, so must do an angular analysis to separate the CP+ and CP- components

# J/ψφ: Transversity

$$\frac{d^4\Gamma(B_s^0 \rightarrow J/\psi\phi)}{dt d\cos\theta d\varphi d\cos\psi} \equiv \frac{d^4\Gamma}{dt d\Omega} \propto \sum_{k=1}^{10} h_k(t) f_k(\Omega)$$

$k$	$h_k(t)$	$f_k(\theta, \psi, \varphi)$
1	$ A_0 ^2(t)$	$2 \cos^2 \psi (1 - \sin^2 \theta \cos^2 \phi)$
2	$ A_{\parallel}(t) ^2$	$\sin^2 \psi (1 - \sin^2 \theta \sin^2 \phi)$
3	$ A_{\perp}(t) ^2$	$\sin^2 \psi \sin^2 \theta$
4	$\Im(A_{\parallel}(t) A_{\perp}(t))$	$-\sin^2 \psi \sin 2\theta \sin \phi$
5	$\Re(A_0(t) A_{\parallel}(t))$	$\frac{1}{2}\sqrt{2} \sin 2\psi \sin^2 \theta \sin 2\phi$
6	$\Im(A_0(t) A_{\perp}(t))$	$\frac{1}{2}\sqrt{2} \sin 2\psi \sin 2\theta \cos \phi$
7	$ A_s(t) ^2$	$\frac{2}{3}(1 - \sin^2 \theta \cos^2 \phi)$
8	$\Re(A_s^*(t) A_{\parallel}(t))$	$\frac{1}{3}\sqrt{6} \sin \psi \sin^2 \theta \sin 2\phi$
9	$\Im(A_s^*(t) A_{\perp}(t))$	$\frac{1}{3}\sqrt{6} \sin \psi \sin 2\theta \cos \phi$
10	$\Re(A_s^*(t) A_0(t))$	$\frac{4}{3}\sqrt{3} \cos \psi (1 - \sin^2 \theta \cos^2 \phi)$



for S-wave under  $\phi$  predicted  
by Stone & Zhang PRD 79,  
074024 (2009)

# Transversity II

$$|A_0|^2(t) = |A_0|^2 e^{-\Gamma_s t} \left[ \cosh\left(\frac{\Delta\Gamma}{2}t\right) - \cos\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) + \sin\phi_s \sin(\Delta m t) \right],$$

$$|A_{\parallel}(t)|^2 = |A_{\parallel}|^2 e^{-\Gamma_s t} \left[ \cosh\left(\frac{\Delta\Gamma}{2}t\right) - \cos\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) + \sin\phi_s \sin(\Delta m t) \right],$$

$$|A_{\perp}(t)|^2 = |A_{\perp}|^2 e^{-\Gamma_s t} \left[ \cosh\left(\frac{\Delta\Gamma}{2}t\right) + \cos\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) - \sin\phi_s \sin(\Delta m t) \right],$$

$$\begin{aligned} \Im(A_{\parallel}^*(t) A_{\perp}(t)) &= |A_{\parallel}| |A_{\perp}| e^{-\Gamma_s t} \left[ -\cos(\delta_{\perp} - \delta_{\parallel}) \sin\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) \right. \\ &\quad \left. - \cos(\delta_{\perp} - \delta_{\parallel}) \cos\phi_s \sin(\Delta m t) + \sin(\delta_{\perp} - \delta_{\parallel}) \cos(\Delta m t) \right], \end{aligned}$$

$$\begin{aligned} \Re(A_0^*(t) A_{\parallel}(t)) &= |A_0| |A_{\parallel}| e^{-\Gamma_s t} \cos(\delta_{\parallel} - \delta_0) \left[ \cosh\left(\frac{\Delta\Gamma}{2}t\right) - \cos\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) \right. \\ &\quad \left. + \sin\phi_s \sin(\Delta m t) \right], \end{aligned}$$

$$\begin{aligned} \Im(A_0^*(t) A_{\perp}(t)) &= |A_0| |A_{\perp}| e^{-\Gamma_s t} \left[ -\cos(\delta_{\perp} - \delta_0) \sin\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) \right. \\ &\quad \left. - \cos(\delta_{\perp} - \delta_0) \cos\phi_s \sin(\Delta m t) + \sin(\delta_{\perp} - \delta_0) \cos(\Delta m t) \right], \end{aligned}$$

$$|A_s(t)|^2 = |A_s|^2 e^{-\Gamma_s t} \left[ \cosh\left(\frac{\Delta\Gamma}{2}t\right) + \cos\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) - \sin\phi_s \sin(\Delta m t) \right], \quad \text{only term for } f=f_{cp}$$

$$\begin{aligned} \Re(A_s^*(t) A_{\parallel}(t)) &= |A_s| |A_{\parallel}| e^{-\Gamma_s t} \left[ -\sin(\delta_{\parallel} - \delta_s) \sin\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) - \sin(\delta_{\parallel} - \delta_s) \cos\phi_s \sin(\Delta m t) \right. \\ &\quad \left. + \cos(\delta_{\parallel} - \delta_s) \cos(\Delta m t) \right], \end{aligned}$$

$$\begin{aligned} \Im(A_s^*(t) A_{\perp}(t)) &= |A_s| |A_{\perp}| e^{-\Gamma_s t} \sin(\delta_{\perp} - \delta_s) \left[ \cosh\left(\frac{\Delta\Gamma}{2}t\right) + \cos\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) \right. \\ &\quad \left. - \sin\phi_s \sin(\Delta m t) \right], \end{aligned}$$

$$\begin{aligned} \Re(A_s^*(t) A_0(t)) &= |A_s| |A_0| e^{-\Gamma_s t} \left[ -\sin(\delta_0 - \delta_s) \sin\phi_s \sinh\left(\frac{\Delta\Gamma}{2}t\right) \right. \\ &\quad \left. - \sin(\delta_0 - \delta_s) \cos\phi_s \sin(\Delta m t) + \cos(\delta_0 - \delta_s) \cos(\Delta m t) \right]. \end{aligned}$$




# Time Dependent CPV in $B_s$ decay

---

- For  $B^0$   $\Delta M = 0.507 \text{ ps}^{-1}$
- For  $B_s$   $\Delta M_s = 17.77 \text{ ps}^{-1}$
- $\therefore$  excellent decay time resolution is required, large  $B_s$  production is also necessary
- Province of hadron collider experiments
- Hints from CDF & D0 for new physics
- New more precise results from LHCb

# LHCb experiment




# The LHCb Collaboration

- 800 Physicists
- 54 Institutes
- 15 Countries
  - 3 Groups from USA




- Basking in light of 2008 Nobel Prize to Kobayashi & Maskawa, “for the discovery of the origin of the broken symmetry which predicts the existence of at least 3 families of quarks”

# LHCb Detector






# Detector Workings





LHCb detector ~ fully installed and commissioned → walk through the detector using the example of a  $B_s \rightarrow D_s K$  decay

# B-Vertex Measurement



Example:  $B_s \rightarrow D_s K$



-5m

## Vertex Locator (Velo)

Silicon strip detector with  
 $\sim 5\text{ }\mu\text{m}$  hit resolution  
 $\rightarrow 30\text{ }\mu\text{m}$  IP resolution


## Vertexing:

- trigger on impact parameter
- measurement of decay distance (time)




# Momentum and Mass measurement


Momentum meas. + direction (VELO):  
Mass resolution for background suppression



Mass resolution  
 $\sigma \sim 15 \text{ MeV}$



$B_s^0 \rightarrow D_s^- K^+$




Sept., 2012



# Hadron Identification

RICH: K/ $\pi$  identification using Cherenkov light emission angle





RICH1: 5 cm aerogel  $n=1.03$   
4 m<sup>3</sup> C<sub>4</sub>F<sub>10</sub>  $n=1.0014$

RICH2: 100 m<sup>3</sup> CF<sub>4</sub>  $n=1.0005$




# Particle identification and L0 trigger





ECAL (inner modules):  $\sigma(E)/E \sim 8.2\% / \sqrt{E} + 0.9\%$

## Calorimeter system :

- Identify electrons, hadrons,  $\pi^0$ ,  $\gamma$
- Level 0 trigger: high  $E_T$  electron and hadron




# Particle identification and L0 trigger




## Muon system:

- Level 0 trigger: High  $P_t$  muons
- OS flavour tagging



# Triggering

**Trigger is crucial as  $\sigma_{b\bar{b}}$  is less than 1% of total inelastic cross section and  $B$  decays of interest typically have  $\mathcal{B} < 10^{-5}$**



□ **Hardware level (L0)**  
Search for high- $p_T$   $\mu$ ,  $e$ ,  $\gamma$  and hadron candidates


□ **Software level (High Level Trigger, HLT)**  
Farm with  $\mathcal{O}(2000)$  multi-core processors  
*HLT1: Confirm L0 candidate with more complete info, add impact parameter and lifetime cuts*  
*HLT2:  $B$  reconstruction + selections*

	$\epsilon(\text{L0})$	$\epsilon(\text{HLT1})$	$\epsilon(\text{HLT2})$
Electromagnetic	70 %	> $\sim 80$ %	> $\sim 90$ %
Hadronic	50 %		
Muon	90 %		



# Running Conditions



VELO rz view



- 20 MHz of bunch crossing (in 2012, with 50 ns bunch spacing) with an average of 2 p-p interactions per bunch crossing → this level of pileup not an issue for LHCb


# Luminosity Leveling

- Luminosity is maintained as at a constant value of  $\sim 4 \times 10^{32} / \text{cm} \cdot \text{s}$  by displacing beams transversely
- Integral  $\mathcal{L}$  is 1/fb in 2011, expect 2.2/fb more in 2012



# $\phi_s$ from $B_s \rightarrow J/\psi \pi^+ \pi^-$

- Reconstructed  $\pi^+ \pi^-$  mass spectrum
- In region between arrows, measured to be  $>97.7\%$  CP-odd @95% cl



- $a[f(t)] \sim 2 \sin \phi_s \sin(\Delta M t)$
- $\phi_s = -0.019^{+0.173+0.004}_{-0.174-0.003} \text{ rad}$

# $\phi_s$ results from $J/\psi\phi$


LHCb values

$$\Gamma = 0.6580 \pm 0.0054 \pm 0.0066 \text{ (ps}^{-1}\text{)}$$

$$\Delta\Gamma = 0.116 \pm 0.018 \pm 0.006 \text{ (ps}^{-1}\text{)}$$

$$\phi_s = 0.001 \pm 0.101 \pm 0.027 \text{ (rad)}$$


Ambiguity removed using interference with  $K^+K^-$  S-wave



- Combining LHCb results:  $\phi_s = -0.002 \pm 0.083 \pm 0.027 \text{ rad}$

# $\Gamma_s$ & $\Delta\Gamma_s$

- $B_s$  lifetime results here use only fully reconstructed decays
- $K^+K^-$  is taken as CP even ( $A_{\Delta\Gamma}=-1$ )
- Ovals show 39% cl, while bands 68% cl
- $\tau_s = 1.509 \pm 0.010$  ps,  
 $\Delta\Gamma_s = 0.092 \pm 0.011$  ps<sup>-1</sup>,  $y_s = \Delta\Gamma_s / 2\Gamma_s = 0.07 \pm 0.01$  (from Anna Phan)





- By definition

$$a_{sl} = \frac{\Gamma(\bar{M} \rightarrow f) - \Gamma(M \rightarrow \bar{f})}{\Gamma(\bar{M} \rightarrow f) + \Gamma(M \rightarrow \bar{f})}$$

at  $t=0$   $\bar{M} \rightarrow f$  is zero as is  $M \rightarrow \bar{f}$

- Here  $f$  is by construction flavor specific,  $f \neq \bar{f}$
- Can measure eg.  $\bar{B}_s \rightarrow D_s^+ \mu^- \nu$ , versus  $B_s \rightarrow D_s^- \mu^+ \nu$ ,
- Or can consider that muons from two B decays can be like-sign when one mixes and the other decays, so look at  $\mu^+ \mu^+$  vs  $\mu^- \mu^-$
- $a_{sl}$  is expected to be very small in the SM,  
 $a_{sl} = (\Delta\Gamma/\Delta M) \tan\phi_{12}$ , where  $\tan\phi_{12} = \text{Arg}(-\Gamma_{12}/M_{12})$
- In SM ( $B^0$ )  $a_{sl}^d = -4.1 \times 10^{-4}$ , ( $B_s$ )  $a_{sl}^s = +1.9 \times 10^{-5}$


# $D^0 a_{sl}$

- Using dimuons ( $3.9\sigma$ )

$$A_{sl}^b = (-0.787 \pm 0.172 \pm 0.093)\%$$



- Indication from D0 that its  $B_s$
- Separate dimuons into  $B_d$  and  $B_s$  samples using muon impact parameter

- Find  $a_{sl}^d = (-0.12 \pm 0.52)\%$   
 $a_{sl}^s = (-1.81 \pm 1.06)\%$



# New D0 Analysis


- Measure  $a_{\text{sl}}^{\text{s}}$  using  $D_s \mu^- \nu$  events,  $D_s \rightarrow \phi \pi^\pm$
- Detect a  $\mu$  associated with a  $D_s$  decay



- Find  $a_{\text{sl}}^{\text{s}} = (-1.08 \pm 0.72 \pm 0.17)\%$
- Also measure  $a_{\text{sl}}^{\text{d}}$  using  $D^+ \mu^- \nu$ ,  $D^+ \rightarrow K \pi^+ \pi^+$
- $a_{\text{sl}}^{\text{d}} = (0.93 \pm 0.45 \pm 0.14)\%$


# $a_{sl}$ according to D0

- $a_{sl}^s = (-1.81 \pm 0.56)\%$
- $a_{sl}^d = (-0.22 \pm 0.30)\%$
- $3\sigma$  from SM
- [arXiv:1208.5813](https://arxiv.org/abs/1208.5813)



# LHCb measurement

- Use  $D_s \mu^- \nu$ ,  $D_s \rightarrow \phi \pi^\pm$ , magnet is periodically reversed. For magnet down:



- Effect of  $B_s$  production asymmetry is reduced to a negligible level by rapid mixing oscillations
- Calibration samples ( $J/\psi$ ,  $D^{*+}$ ) used to measure detector trigger, track & muon ID biases

# $a_{sl}$ not D0

- LHCb finds


$$a_{sl}^s = (-0.24 \pm 0.54 \pm 0.33)\%$$

- B-factory

$$a_{sl}^d = (-0.05 \pm 0.56)\%$$

- Results consistent with SM

- Expect  $\phi_s$  to grow as  $\sin[2|\beta_s| + \arg(M_{12}^s)]$  for finite  $a_{sl}$ .




# CPV in Charm

- Expect largest effects in Cabibbo Suppressed Decays. COULD REVEAL NP (see Grossman Kagan & Nir [arXiv:1204.3557](https://arxiv.org/abs/1204.3557))
- Define:  $A_{CP}(D \rightarrow f) = \frac{\Gamma(D \rightarrow f) - \Gamma(\bar{D} \rightarrow \bar{f})}{\Gamma(D \rightarrow f) + \Gamma(\bar{D} \rightarrow \bar{f})}$ , if  $f$  is a CP eigenstate then  $f = \bar{f}$
- Current data mainly from LHCb, CDF & Belle show
 
$$\Delta A_{CP} \equiv A_{CP}(K^+ K^-) - A_{CP}(\pi^+ \pi^-) = (-0.74 \pm 0.15)\%$$
- A 4.5  $\sigma$  effect (|| talks Tico, Tonelli) & Ko
- Both SM & NP explanations are prolific
- Choose to treat this as a limit on NP:  $1\% > -\Delta A_{CP} > 0\%$

# $B \rightarrow K^{(*)} e^+ e^-$

- Similar to  $K^* \gamma$ , but more decay paths




+ new particles in loops

- Several variables can be examined, e.g. muon forward-backward asymmetry,  $A_{FB}$  is well predicted in SM




# $B^0 \rightarrow K^{*0} \ell^+ \ell^-$



LHCb ||  
Gallas  
Torreira,  
BaBar ||  
Eigen,  
CDF ||  
Miyake,  
Belle  
PRL 103,  
171801  
(2009)

# Forward-Backward asymmetry






No evidence of deviation from SM so far



# Isospin asymmetry

$$\frac{\Gamma(B^0 \rightarrow K^{(*)0} \mu^+ \mu^-) - \Gamma(B^+ \rightarrow K^{(*)+} \mu^+)}{\Gamma(B^0 \rightarrow K^{(*)0} \mu^+ \mu^-) + \Gamma(B^+ \rightarrow K^{(*)+} \mu^+)}$$



Not SM, but no NP model yet.  
Annihilation diagram only for  $B^-$ , but why the difference for  $K^*$  &  $K$ ?

# Other Processes

- Other processes probe different operators

- Time dependent CPV in  $B^0 \rightarrow K^* \gamma$ ,  $K^* \rightarrow K_S \pi^0$ , is given by

$$\frac{\Gamma(\bar{B}^0(t) \rightarrow \bar{K}^{*0} \gamma) - \Gamma(B^0(t) \rightarrow K^{*0} \gamma)}{\Gamma(\bar{B}^0(t) \rightarrow \bar{K}^{*0} \gamma) + \Gamma(B^0(t) \rightarrow K^{*0} \gamma)} = S_{K^* \gamma} \sin(\Delta M_d t) - C_{K^* \gamma} \cos(\Delta M_d t)$$

where  $S_{K^* \gamma} = -2.3\%$  in SM

- For Generic NP

$$S_{K^* \gamma} \simeq \frac{2}{|C_7|^2 + |C'_7|^2} \text{Im}(e^{-2i\beta} C_7 C'_7)$$

- Data, BaBar & Belle  $(-16 \pm 22)\%$ , still useful even with the large error

# Rare Decays - Generic

- $$\mathcal{H}_{\text{eff}} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \frac{e^2}{16\pi^2} \sum_i (C_i O_i + C'_i O'_i) + \text{h.c.} .$$

- $C_i O_i$  for SM,  $C'_i O'_i$  are for NP. Operators are for  $P_{R,L} = (1 \pm \gamma_5)/2$

$$O_7 = \frac{m_b}{e} (\bar{s} \sigma_{\mu\nu} P_R b) F^{\mu\nu}, \quad O_8 = \frac{gm_b}{e^2} (\bar{s} \sigma_{\mu\nu} T^a P_R b) G^{\mu\nu a},$$


$$O_9 = (\bar{s} \gamma_\mu P_L b) (\bar{\ell} \gamma^\mu \ell), \quad O_{10} = (\bar{s} \gamma_\mu P_L b) (\bar{\ell} \gamma^\mu \gamma_5 \ell),$$

$$O_S = m_b (\bar{s} P_R b) (\bar{\ell} \ell), \quad O_P = m_b (\bar{s} P_R b) (\bar{\ell} \gamma_5 \ell),$$

- $O' = O$  with  $P_{R,L} \rightarrow P_{L,R}$
- Each process depends on a unique combination

# Common Analysis

- APS  $\equiv$  W. Altmannshofer, P. Paradisi & D. M. Straub arXiv:1111.1257v2



- Many more such generic constraints




# $B_s \rightarrow \mu^+ \mu^-$


- SM branching ratio is  $(3.2 \pm 0.2) \times 10^{-9}$  [Buras arXiv: 1012.1447], NP can make large contributions.

Note, K. De Brun arXiv:1204.1737 show that B theory needs to be raised by  $1/(1-y_s)$

Standard Model





MSSM



- Many NP models possible, not just Super-Sym

# Discrimination

- LHCb & CDF use  $B \rightarrow h^+ h^-$  to tune cuts. They use a multivariate analysis
- Other variables to discriminate against bkgd : B impact parameter, B lifetime, B  $p_t$ , B isolation, muon isolation, minimum impact parameter of muons, ...
- CMS & ATLAS use  $f_s/f_d$  from LHCb





See ICHEP talk of M. Perrin-Terrin

# ATLAS+CMS+LHCb


- CLs for bkgnd only, dashed line is the expectation, blue curve show the measurement, red the 95% cl limit
- LHCb data show slight excess consistent with SM
- Also

$$\mathcal{B}(B_d \rightarrow \mu^+ \mu^-) < 8.1 \times 10^{-10}$$



# Results



95% confidence level limits



# Implications


- “LHC” limit
    - $<4.2 \times 10^{-9}$  @95% CL
    - This is 1.2 times SM value
  - Set serious limits in NUHM1 SUSY model
  - Other LHCb results
    - $\mathcal{B}(B_s \rightarrow \mu^+ \mu^- \mu^+ \mu^-) < 1.3 \times 10^{-8}$
    - $\mathcal{B}(B_d \rightarrow \mu^+ \mu^- \mu^+ \mu^-) < 5.4 \times 10^{-9}$
- Predicted via “portals”

see arXiv:0911.4938



# Implications II

David Straub, Rencontres de Moriond EW, La Thuile (2012)



The 125 GeV Higgs observations kills off 4<sup>th</sup> generation models as the production cross-section would be 9x larger & decays to  $\gamma\gamma$  suppressed


Sept., 2012





# $B^- \rightarrow \tau^- \bar{\nu}$ problem?

- $B^- \rightarrow \tau^- \bar{\nu}$ , tree process:




Can be new particles instead of  $W^-$  but why not also in  $D_{(s)}^+ \rightarrow \ell^+ \nu$ ?

- $\sin 2\beta$ , CPV in e.g.  $B^0 \rightarrow J/\psi K_s$ : Box diagram


- Measurement not in good agreement with SM prediction based on CKM fit (Yook ICHEP talk)

Discrepancy may be resolved; what caused the change?



# Peaking Backgrounds


- Since  $e^+e^- \rightarrow B^+B^-$ , analysis uses reconstruction of  $B^+$ , detection of  $\tau^- \rightarrow$  one track & small extra E



# $B \rightarrow D^{(*)} \tau \nu$

- Also, tree level –new BaBar result

- Similar to  $B^- \rightarrow \tau^- \nu$  analysis: fully reconstruct one B, keep events with an additional  $D^{(*)}$  plus an  $e^-$  or  $\mu^-$ .




- Signal is wide, background, especially  $D^{**} \ell \nu$ , needs careful estimation

# BaBar results

- Results given in terms of ratio to  $B \rightarrow D^{(*)} \ell \nu$


	SM Theory	BaBar value	Diff.
$R(D)$	$0.297 \pm 0.017$	$0.440 \pm 0.058 \pm 0.042$	$+2.0\sigma$
$R(D^*)$	$0.252 \pm 0.003$	$0.332 \pm 0.024 \pm 0.018$	$+2.7\sigma$

- Sum is  $3.4\sigma$  above SM
- Also inconsistent with type II 2HDM  
(see De Nardo ICHEP talk)



# Belle Results

- Two types of analysis, hadronic tags (arXiv: 0910.4301) similar to BaBar and also “inclusive tags” (A. Matyja et. al, PRL 99,191807 (2007)).



- Belle data currently support BaBar indication of larger than expected rates
- Belle should be able to reduce uncertainties to the BaBar level
- Will be interesting to see results of 2D fits


# The Dark Sector

- Could it be that there are 3 classes of matter?
  - SM particles with charges  $[SU(3) \times SU(2) \times U(1)]$
  - Dark matter particles with “dark” charges
  - Some matter having both (“mediators”)
- Searches for “dark photons”
  - A mediator, couples to b-quarks (see arXiv:056151 hep/ph)
  - BaBar  $\mathcal{B}(Y(1S) \rightarrow \text{invisible}) < 3 \times 10^{-4}$  @ 90% cl
  - Other experiments




# Search Summary

- Parameterize by mixing  $\epsilon$



- Dark photon mass  $m_{A'}$




From B. Echenard arXiv:1205.3505

# Dark Higgs



- BaBar search for  $e^+e^- \rightarrow h'A'$ ,  $h' \rightarrow A'A'$
- $A'$  is looked for in  $e^+e^-$ ,  $\mu^+\mu^-$ ,  $\tau^+\tau^-$  & hadrons
- Limits parameterized in terms of mixing  $\varepsilon$  & dark matter coupling  $\alpha_D$

- Nothing found, upper limits set at 90% cl:



# Majorana $\nu$ 's

- Several ways of looking for presence of heavy  $\nu$ 's (N) in heavy quark decays if they Majorana (their own anti-particles) and couple to "ordinary"  $\nu$ 's
- Modes analogous to  $\nu$ -less nuclear  $\beta$  decay



Simplest Channels:

$$B^- \rightarrow D^+ \ell^- \ell'^- \text{ \& } B^- \rightarrow D^{*+} \ell^- \ell'^-$$

$$B^- \rightarrow D^{*+} \ell^- \ell'^-$$

$\ell^-$  &  $\ell'^-$  can be  $e^-$ ,  $\mu^-$  or  $\tau^-$ .

# Limits on $D^{(*)+} \ell^- \ell'^-$


- Upper limits in  $e^-e^-$  mode not competitive with nuclear  $\beta$  decay
- Others unique since measure coupling of Majorana  $\nu$  to  $\mu^-$

Mode	Exp.	u. l. $\times 10^{-6}$
$B^- \rightarrow D^+ e^- e^-$	Belle	$< 2.6$
$B^- \rightarrow D^+ e^- \mu^-$	Belle	$< 1.8$
$B^- \rightarrow D^+ \mu^- \mu^-$	Belle	$< 1.0$
$B^- \rightarrow D^+ \mu^- \mu^-$	LHCb	$< 0.69$
$B^- \rightarrow D^{*+} \mu^- \mu^-$	LHCb	$< 3.6$


Belle [arXiv:1107.064]

# On-Shell $\nu$

- Can also look for Majorana  $\nu$  (N), where  $N \rightarrow W^+ \mu^-$
- Several ways
- A. Atre, T. Han, S. Pascoli, & B. Zhang [arXiv:0901.3589]
- N. Quintero, G. Lopez & Castro, [arXiv:1108.6009]






# LHCb searches



Nothing  
yet

Aaij, PRD 85,  
112004 (2012)





# Conclusions

- Although there is no compelling evidence yet for NP, Flavor physics is very sensitive to potential effects at high mass scales. All NP theories must satisfy stringent experimental constraints
- Experiments have been very effective at dispelling effects with marginal statistical significance, although a few remain. Will some stand when precision improves?
- Improving measurements such as  $B_s \rightarrow \mu^+ \mu^-$ ,  $B \rightarrow K \mu^+ \mu^-$ , CPV:  $\phi_s$ , etc., may show NP effects, & need to be aggressively pursued
- We are looking forward to new flavor physics discoveries from the LHC & its upgrades, BESIII, and Super B factories
- We are looking forward to defining the next theory beyond the SM

# Theory conquers



---

# *The End*

---